

Calculation of entanglement for continuous variable states

Huai-Xin Lu,^{1,2} Zeng-Bing Chen,¹ Jian-Wei Pan,^{1,3} and Yong-De Zhang^{4,5}

¹Department of Modern Physics, University of Science and Technology of China, Hefei, Anhui 230027, China ²Department of Physics, Wefang University, Wefang, Shandong 261043, China

³Institut für Experimentalphysik, Universität Wien, Boltzmanngasse 5, 1090 Wien, Austria

⁴CCAST (World Laboratory), P.O. Box 8730, Beijing 100080, China

⁵Laboratory of Quantum Communication and Quantum Computation and Department of Modern Physics, University of Science and Technology of China, Hefei, Anhui 230027, China

()

In this paper, we present a general formula for obtaining the reduced density operator for any biparticle pure entangled state. Using this formula, we derive, in a compact form, the explicit formula of the entanglement for any bipartical pure entangled Gaussian state. In the case of Gaussian states, the criteria of separability can be naturely obtained by the formula. For non-Gaussian states, we also show the usefulness of the method presented in this paper.

PACS: 03.65.Ud, 42.50.-p

Quantum entanglement is one of the essential features of quantum mechanics. It has been an interesting and important topic since it was first noted by Einstein, Podolsky, and Rosen (EPR) [1]. Now, it has attracted much attention because of immense progresses both in the foundation of quantum mechanics and in the burgeoning field of quantum information theory. Issues such as the relationship between entanglement and quantum nonlocality and the violations of Bell's inequality for both discrete [2,3] and continuous variable states [4] further enrich the contents of quantum mechanics. In recent years, quantum entanglement is also viewed as a useful "resource" for various kinds of quantum information processing. The original EPR states of continuous variables can be created either by the nondegenerate optical parametric amplifier [5–7], or simply by using passive optical elements (e.g., beam splitters) [8–10]. While most quantum information protocols were initially developed for quantum systems with discrete quantum variables, quantum information processing based on quantum states with continuous variable has also been proposed, e.g., quantum teleportation [11], quantum error correction [12], quantum computation [13], entanglement purification [14] and cloning [15]. Thus, inseparability criteria [16–18] and entanglement quantification [19] in continuous variable systems has become an issue of practical importance.

The measurement of entanglement for any biparticle pure entangled state, i.e., the von Neumann entropy of either partial trace of the density operator for the state, is considered as a "good measurement" of quantum entanglement. Parker *et al.* [19] have developed an elegant method to calculate the entanglement of the biparticle pure entangled states with continuous variables by means

of the integral eigenvalue equations in the coordinate-momentum space. In fact, their method is based on the Schmidt decompositions. But in certain cases, it is somewhat harder to find such a "Schmidt basis" for the continuous variable systems. In this paper, we propose an alternative method to calculate the entanglement entropy for continuous variable states in Fock space. With the help of linear quantum transformation theory (LQTT) [20], we obtain an explicit formula of the entanglement entropy for *any* Gaussian state. With this formula, we can easily calculate the Gaussian states' entanglement without knowing the eigenvalues of the reduced density operator for the system. Furthermore, from the derivation of the formula, we naturally obtain the necessary and sufficient condition of separability for a biparticle pure entangled state with continuous variables. For some non-Gaussian states, one can also calculate the eigenvalues of the reduced density operators, from which the entanglement of this bipartite system may be easily obtained.

It is well known that the entanglement entropy of a biparticle pure entangled state is defined by

$$E = -\text{tr } \rho_1 \ln \rho_1 = -\text{tr } \rho_2 \ln \rho_2, \quad (1)$$

where the reduced density operator $\rho_{1(2)}$ is a partial trace of density operator ρ_{12} for the state, i.e.,

$$\rho_{1(2)} = \text{tr}_{2(1)} \rho_{12}. \quad (2)$$

In order to give a general formula of reduced density operator ρ_1 for any biparticle system, we use the following formula [20,21]

$$\Omega = : \exp \left[(a_1^\dagger, a_2^\dagger) \begin{pmatrix} \partial_{z_1^*} \\ \partial_{z_2^*} \end{pmatrix} + (\partial_{z_1}, \partial_{z_2}) \begin{pmatrix} a_1 \\ a_2 \end{pmatrix} \right] : \\ \times \langle Z | \Omega | Z \rangle |_{z=0}, \quad (3)$$

where Ω is an arbitrary bosonic operator, $a_i^\dagger$ ($i = 1, 2$) is the usual bosonic creation operator for the subsystem i ; $|Z\rangle = |Z_1\rangle \otimes |Z_2\rangle$ denote the usual coherent states, $::$ means the normal ordering and $z = 0$ means $z_{1(2)} = z_{1(2)}^* = 0$. Using Eq. (3), we write the density operator ρ_{12} for the biparticle system as

$$\rho_{12} = : \exp \left[(a_1^\dagger, a_2^\dagger) \begin{pmatrix} \partial_{z_1^*} \\ \partial_{z_2^*} \end{pmatrix} + (\partial_{z_1}, \partial_{z_2}) \begin{pmatrix} a_1 \\ a_2 \end{pmatrix} \right] : \\ \times \langle Z | \rho_{12} | Z \rangle |_{z=0}, \quad (4)$$

where $|Z\rangle \equiv |Z_1, Z_2\rangle$. With the help of overcomplete of coherent states $\int \frac{d^2 Z_2}{\pi} |Z_2\rangle_{22} \langle Z_2| = 1$, and taking the partial trace over subspace of the subsystem 2, one can get the reduced density operator ρ_1 as follows

$$\begin{aligned} \rho_1 &= \text{tr}_2 \rho_{12} = \text{tr}_2 \int \frac{d^2 z_2'}{\pi} |Z'\rangle_{22} \langle Z'| \\ &\quad \times : e^{(a_1^\dagger, a_2^\dagger) \begin{pmatrix} \partial_{z_1'}^* \\ \partial_{z_2'}^* \end{pmatrix} + (\partial_{z_1}, \partial_{z_2}) \begin{pmatrix} a_1 \\ a_2 \end{pmatrix}} : \langle Z | \rho_{12} | Z \rangle |_{z=0} \\ &= \int \frac{d^2 z_2'}{\pi} : \exp \left(a_1^\dagger \partial_{z_1'} + \partial_{z_1} a_1 \right) : \\ &\quad \times \exp \left(z_2'^* \partial_{z_2'} + \partial_{z_2} z_2'^* \right) \langle Z | \rho_{12} | Z \rangle |_{z=0}. \end{aligned} \quad (5)$$

In the following we will consider two classes of states: the Gaussian states and the non-Gaussian ones. Let us first consider the following Gaussian states

$$\begin{aligned} \rho_{12} &= A_0 : \exp \left\{ \frac{1}{2} \left[(a_1^\dagger, a_1) M_1 \begin{pmatrix} a_1^\dagger \\ a_1 \end{pmatrix} \right. \right. \\ &\quad \left. \left. + (a_2^\dagger, a_2) M_2 \begin{pmatrix} a_2^\dagger \\ a_2 \end{pmatrix} + 2(a_1^\dagger, a_1) M_{12} \begin{pmatrix} a_2^\dagger \\ a_2 \end{pmatrix} \right] \right\} : , \end{aligned} \quad (6)$$

where A_0 is a normalization factor, M_i ($i = 1, 2$) is Hermitian matrices, and

$$M_{12} = \begin{pmatrix} e & f \\ f^* & e^* \end{pmatrix}$$

with e and f being two arbitrary complex numbers. In Eq. (6) the linear terms of a and $a^\dagger$ are not included since they do not affect the entanglement of the bipartite Gaussian states [16]. In this case, the matrix elements of ρ_{12} with respect to $|Z\rangle$ is

$$\begin{aligned} \langle Z | \rho_{12} | Z \rangle &= A_0 \exp \left\{ \frac{1}{2} \left[(z_1^*, z_1) M_1 \begin{pmatrix} z_1^* \\ z_1 \end{pmatrix} + (z_2^*, z_2) \right. \right. \\ &\quad \left. \left. \times M_2 \begin{pmatrix} z_2^* \\ z_2 \end{pmatrix} + 2(z_1^*, z_1) M_{12} \begin{pmatrix} z_2^* \\ z_2 \end{pmatrix} \right] \right\}. \end{aligned} \quad (7)$$

Using the following Gaussian integration formula [22]

$$\begin{aligned} &\int \frac{d^2 z}{\pi} \exp \left\{ -\frac{1}{2} (z^*, z) Q \begin{pmatrix} z^* \\ z \end{pmatrix} + (u, v) \begin{pmatrix} z^* \\ z \end{pmatrix} \right\} \\ &= [-\det Q]^{\frac{-1}{2}} \exp \left\{ \frac{1}{2} (u, v) Q^{-1} \begin{pmatrix} u \\ v \end{pmatrix} \right\}, \end{aligned} \quad (8)$$

where $Q = \tilde{Q}$ is nonsingular and u (v) is an arbitrary complex number, and the formula

$$\begin{aligned} &\exp \left(z_2'^* \partial_{z_2'} + \partial_{z_2} z_2'^* \right) \Psi(z_1^*, z_1; z_2^*, z_2) |_{z_2=z_2'=0} \\ &= \Psi(z_1^*, z_1; z_2'^*, z_2'), \end{aligned} \quad (9)$$

we have, after substituting Eq. (7) into Eq. (5),

$$\rho_1 = \frac{A_0}{\sqrt{-\det M_2}} : e^{\frac{1}{2} (a_1^\dagger, a_1) (M_1 - M_{12} M_2^{-1} \tilde{M}_{12}) \begin{pmatrix} a_1^\dagger \\ a_1 \end{pmatrix}} : . \quad (10)$$

According to LQTT [20], if one denotes

$$M = \begin{pmatrix} a & d \\ b & c \end{pmatrix}, \quad (11)$$

there is a map of M defined as

$$\begin{aligned} D(M) &= (M_1 - M_{12} M_2^{-1} \tilde{M}_{12}) \Sigma_B^{-1} \\ &= \begin{pmatrix} c^{-1} - 1 & c^{-1} d \\ c^{-1} b & 1 - c^{-1} \end{pmatrix} \Sigma_B^{-1}, \end{aligned} \quad (12)$$

where complex constants a, b, c and d are determined by the known matrices $M_{1(2)}$ and M_{12} in Eq. (10). Then we can immediately rewrite the normally ordered form in Eq. (10) as [20]

$$\rho_1 = A \exp \left[\frac{1}{2} (a_1^\dagger, a_1) N \Sigma_B \begin{pmatrix} a_1^\dagger \\ a_1 \end{pmatrix} \right], \quad (13)$$

where negative Hermitian matrix $N = \ln M$, $\Sigma_B = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$, and A is constant

$$A = A_0 \sqrt{\frac{c}{-\det M_2}}. \quad (14)$$

Substituting Eq. (13) into Eq. (1), we get

$$E = -A \left[Z(\beta) \ln A - \frac{d}{d\beta} Z(\beta) \right]_{\beta=-1}, \quad (15)$$

in which

$$Z(\beta) = \text{tr} \exp \left\{ -\frac{\beta}{2} (a_1^\dagger, a_1) N \Sigma_B \begin{pmatrix} a_1^\dagger \\ a_1 \end{pmatrix} \right\}. \quad (16)$$

Using the result in Ref. [22]

$$\begin{aligned} Z(\beta) &= \text{tr} \exp \left\{ -\frac{\beta}{2} (a_1^\dagger, a_1) N \Sigma_B \begin{pmatrix} a_1^\dagger \\ a_1 \end{pmatrix} \right\} \\ &= |\det(e^{\beta N} - 1)|^{-\frac{1}{2}}, \end{aligned} \quad (17)$$

direct calculation yields

$$\frac{d}{d\beta} Z(\beta) |_{\beta=-1} = -\frac{1}{2} |\det(e^{-N} - 1)|^{-\frac{1}{2}} \text{tr} \frac{N}{1 - e^N}. \quad (18)$$

Substituting Eq. (18) into Eq. (15) we get the explicit expression of the entanglement for the state in Eq. (6)

$$E = -A |\det(e^{\beta N} - 1)|^{-\frac{1}{2}} \left[\ln A + \frac{1}{2} \text{tr} \frac{N}{1 - e^N} \right]. \quad (19)$$

Thus, from Eq. (19) one can conveniently obtain the entanglement of states in Eq. (6) only by calculating the

eigenvalues of negative Hermitian matrix N , which can be easily obtained. This is the main result of this paper. It is worth pointing out that from Eq. (19), the necessary and sufficient condition of separability for the states in Eq. (6) naturally reads

$$\ln A = \frac{1}{2} \text{tr} \frac{N}{e^N - 1}. \quad (20)$$

As an application of Eq. (19), we consider the entangled states produced by a beam splitter [10]

$$|\psi\rangle_{12} = \hat{B}(\theta, \phi) \hat{S}_1(\zeta_1) \hat{S}_2(\zeta_2) |00\rangle, \quad (21)$$

where the beam splitter operator $\hat{B}(\theta, \phi)$ is [23]

$$\hat{B}(\theta, \phi) = \exp \left[\theta (a_1^\dagger a_2 e^{i\varphi} - a_1 a_2^\dagger e^{-i\varphi}) \right], \quad (22)$$

and the single-mode squeezed operator $\hat{S}(\zeta)$ is [24]

$$\hat{S}(\zeta) = \exp \left[\frac{1}{2} (\zeta^* a^2 - \zeta a^{\dagger 2}) \right]. \quad (23)$$

Making use of LQTT, we can write the normally ordered form of the density operator in Eq. (21) as follows

$$\begin{aligned} \rho_{12} = |\psi\rangle_{1212} \langle\psi| = A_0 : \exp \left\{ \frac{1}{2} \left[(a_1^\dagger, a_1) M_1 \begin{pmatrix} a_1^\dagger \\ a_1 \end{pmatrix} \right. \right. \\ \left. \left. + (a_2^\dagger, a_2) M_2 \begin{pmatrix} a_2^\dagger \\ a_2 \end{pmatrix} + 2(a_1^\dagger, a_1) M_{12} \begin{pmatrix} a_2^\dagger \\ a_2 \end{pmatrix} \right] \right\} : , \quad (24) \end{aligned}$$

where

$$\begin{aligned} M_1 &= - \begin{pmatrix} \alpha & 1 \\ 1 & \alpha^* \end{pmatrix}, \quad M_2 = - \begin{pmatrix} \beta & 1 \\ 1 & \beta^* \end{pmatrix}, \\ M_{12} &= - \begin{pmatrix} \delta & 0 \\ 0 & \delta^* \end{pmatrix}, \\ \alpha &= \frac{\zeta_1}{|\zeta_1|} \tanh |\zeta_1| \cos^2 \theta + e^{2i\varphi} \frac{\zeta_2}{|\zeta_2|} \tanh |\zeta_2| \sin^2 \theta, \\ \beta &= e^{-2i\varphi} \frac{\zeta_1}{|\zeta_1|} \tanh |\zeta_1| \sin^2 \theta + \frac{\zeta_2}{|\zeta_2|} \tanh |\zeta_2| \cos^2 \theta, \\ \delta &= \frac{1}{2} \sin(2\theta) \left(\frac{\zeta_2}{|\zeta_2|} \tanh |\zeta_2| e^{i\varphi} - \frac{\zeta_1}{|\zeta_1|} \tanh |\zeta_1| e^{-i\varphi} \right). \end{aligned}$$

Using Eqs. (14) and (12) yields

$$A = \frac{1}{|\delta| \cosh |\zeta_1| \cosh |\zeta_2|}, \quad M = \begin{pmatrix} a & d \\ b & c \end{pmatrix} \quad (25)$$

with

$$\begin{aligned} a &= \frac{1}{c} (1 + bd), \quad c = \frac{1 - |\beta|^2}{|\delta|^2}, \\ d &= \frac{1}{|\delta|^2} \left(\delta^2 \beta^* + \alpha (1 - |\beta|^2) \right), \quad b = -d^*. \quad (26) \end{aligned}$$

Finally, we get the entanglement of $|\psi\rangle_{AB}$ in Eq. (21)

$$\begin{aligned} E &= \frac{1}{|\delta| \cosh |\zeta_1| \cosh |\zeta_2|} \\ &\times \frac{\sqrt{\lambda}}{|\lambda - 1|} \left[\ln(|\delta| \cosh |\zeta_1| \cosh |\zeta_2|) \right. \\ &\left. - \frac{1 + \lambda}{2(1 - \lambda)} \ln \lambda \right], \quad (27) \end{aligned}$$

where λ is any of the roots in the following equation

$$\lambda^2 - (a + c)\lambda + 1 = 0. \quad (28)$$

We can consider a special case of Eq. (21) by choosing $\theta = \frac{\pi}{4}$, $\zeta_i = s_i e^{i\varphi}$ ($i = 1, 2$) and $\varphi = \frac{l\pi}{2}$ ($l = 0, 1, 2, \dots$) [10]. In this case Eq. (27) is just the result of Ref. [10], i.e.,

$$E = \cosh^2 |s| \ln(\cosh^2 |s|) - \sinh^2 |s| \ln(\sinh^2 |s|). \quad (29)$$

where $s = \frac{1}{2}(s_1 e^{i\varphi} + s_2 e^{-i\varphi})$. Furthermore, if we choose $\varphi = 0$, $\zeta_2 = -\zeta_1 = r$ and $\theta = \frac{\pi}{4}$, we get $\lambda_1 = \tanh^2 r$. we then get the well known result for the two-mode squeezed vacuum state, whose entanglement is [19]

$$E = \cosh^2 r \ln(\cosh^2 r) - \sinh^2 r \ln(\sinh^2 r). \quad (30)$$

For non-Gaussian states, generally speaking, their entanglement can not be written in the compact form as in Eq. (19). But for some states such as $SU(2)$ coherent states (see, e.g., [10]) and the photon-added coherent states [25], the entanglement can be easily obtained by solving the eigenvalue problem of the reduced operator of Eq. (5). To show this, we consider the following example [10]

$$|\psi\rangle = \hat{B} |n_1, n_2\rangle = e^{\theta(a_1^\dagger a_2 - a_2^\dagger a_1)} |n_1, n_2\rangle \quad (31)$$

The density operator ρ_{12} of this system can be written, by following the above procedure, as

$$\begin{aligned} \rho_{12} &= \hat{B} |n_1, n_2\rangle \langle n_2, n_1| \hat{B}^\dagger = \frac{1}{n_1! n_2!} \frac{d^{n_1}}{d\alpha^{n_1}} \frac{d^{n_2}}{d\beta^{n_2}} \\ &\times : \exp \left\{ \frac{1}{2} \left[(a_1^\dagger, a_1) M_1 \begin{pmatrix} a_1^\dagger \\ a_1 \end{pmatrix} + (a_2^\dagger, a_2) M_2 \begin{pmatrix} a_2^\dagger \\ a_2 \end{pmatrix} \right. \right. \\ &\left. \left. + 2(a_1^\dagger, a_1) M_{12} \begin{pmatrix} a_2^\dagger \\ a_2 \end{pmatrix} \right] \right\} :_{|\alpha=\beta=0}. \quad (32) \end{aligned}$$

Here

$$\begin{aligned} M_1 &= (\alpha \cos^2 \theta + \beta \sin^2 \theta - 1) \sigma_1, \\ M_2 &= (\beta \cos^2 \theta + \alpha \sin^2 \theta - 1) \sigma_1, \\ M_3 &= \sin \theta \cos \theta (\beta - \alpha) \sigma_1, \quad (33) \end{aligned}$$

where σ_1 is Pauli matrix. Using Eq. (5), we derive

$$\rho_A = \frac{1}{n_1!n_2!} \frac{d^{n_1}}{d\alpha^{n_1}} \frac{d^{n_2}}{d\beta^{n_2}} \frac{1}{(\beta \cos^2 \theta + \alpha \sin^2 \theta - 1)} \times \left(\frac{\alpha\beta - (\alpha \cos^2 \theta + \beta \sin^2 \theta)}{(\beta \cos^2 \theta + \alpha \sin^2 \theta - 1)} \right)^{a_1^\dagger a_1} \Big|_{\alpha=\beta=0}. \quad (34)$$

Thus we get the eigenvalues as follows

$$\lambda_{N_1} = \frac{1}{n_1!n_2!} \frac{d^{n_1}}{d\alpha^{n_1}} \frac{d^{n_2}}{d\beta^{n_2}} \frac{1}{\beta \cos^2 \theta + \alpha \sin^2 \theta} \times \left[1 + \frac{\alpha\beta - 1}{\beta \cos^2 \theta + \alpha \sin^2 \theta} \right]^{N_1} \Big|_{\alpha=\beta=-1} \quad (35)$$

Note that in Eq. (35) we take the replacements $\alpha = \alpha' + 1$ and $\beta = \beta' + 1$. Finally, we get the entanglement of the state in Eq. (31)

$$E = - \sum_{N_1} \lambda_{N_1} \ln \lambda_{N_1}. \quad (36)$$

In summary, we have presented a general formula of reduced density operator for any biparticle pure entangled state of continuous variables. Based on this formula, we have derived the explicit expression of the entanglement for any Gaussian pure state in a compact form. From the examples mentioned above, it is easy to see that Gaussian states' entanglement can be conveniently calculated by the formula given in this paper. For the case of non-Gaussian states, we have also shown the usefulness of the formula for calculating the reduced density operator, as given above. In this sense, the method to calculate the entanglement for biparticle pure entangled state is easier to operate.

This work was supported by the National Natural Science Foundation of China under Grants No. 19975043, No. 10104014 and No. 10028406, and by the Chinese Academy of Sciences.

- [6] Z. Y. Ou, S. F. Pereira, H. J. Kimble, and K. C. Peng, Phys. Rev. Lett. **68**, 3663 (1992).
- [7] Z. Y. Ou, S. F. Pereira, and H. J. Kimble, Appl. Phys. B: Photophys. Laser Chem. **55**, 265 (1992).
- [8] P. van Loock and S.L. Braunstein, Phys. Rev. Lett. **84**, 3482 (2000).
- [9] S. Scheel and D.-G. Welsch, Phys. Rev. A **64**, 063811 (2001).
- [10] M.S. Kim, W. Son, V. Bužek, and P.L. Knight, *ibid.* **65**, 032323 (2002).
- [11] L. Vaidman, Phys. Rev. A **49**, 1473 (1994); S. L. Braunstein and H. J. Kimble, Phys. Rev. Lett. **80**, 869 (1998); A. Furusawa *et al.*, Science **282**, 706 (1998).
- [12] S. L. Braunstein, Phys. Rev. Lett. **80**, 4084 (1998); S. Lloyd and J.-J. E. Slotine, *ibid.* **80**, 4088 (1998); S. L. Braunstein, Nature (London) **394**, 47 (1998).
- [13] S. Lloyd and S. L. Braunstein, Phys. Rev. Lett. **82**, 1784 (1999).
- [14] L.-M. Duan *et al.*, Phys. Rev. Lett. **84**, 4002 (2000).
- [15] N. J. Cerf, A. Ipe, and X. Rottenberg, Phys. Rev. Lett. **85**, 1754 (2000).
- [16] L.-M. Duan, G. Giedke, J. I. Cirac, and P. Zoller, Phys. Rev. Lett. **84**, 2722 (2000).
- [17] R. Simon, Phys. Rev. Lett. **84**, 2726 (2000).
- [18] Wang Xiang-Bin, M. Keiji, and T. Akihisa, Phys. Rev. Lett. **87**, 137903 (2001).
- [19] S. Parker, S. Bose, and M.B. Plenio, Phys. Rev. A **61**, 032305 (2000).
- [20] Y.-D. Zhang and Z. Tang, J. Math. Phys. **34**, 5639 (1993).
- [21] C. Itzykson and J. B. Zuber, Quantum Field Theory (McGraw-Hill, NY, 1980).
- [22] J.-W. Pan, Q.-X. Dong, Y.-D. Zhang, G. Hou, and X.-B. Wang, Phys. Rev. E **56**, 2553 (1997).
- [23] R. A. Campos, B. E. A. Saleh, and M. C. Teich, Phys. Rev. A **40**, 1371 (1989).
- [24] R. Loudon and P. L. Knight, J. Mod. Opt. **34**, 709 (1987).
- [25] G. S. Agarwal and K. Tara, Phys. Rev. A **43** 492 (1991).

-
- [1] A. Einstein, B. Podolsky and N. Rosen, Phys. Rev. **47**, 777 (1935).
 - [2] J. S. Bell, Physics (Long Island) **1**, 195 (1964); *Speakable and Unsayable in Quantum Mechanics* (Cambridge Univ. Press, Cambridge, 1987).
 - [3] J. F. Clauser, M. A. Horne, A. Shimony and R. A. Holt, Phys. Rev. Lett. **23**, 880 (1969).
 - [4] K. Banaszek and K. Wódkiewicz, Phys. Rev. A **58**, 4345 (1998); Phys. Rev. Lett. **82**, 2009 (1999); Acta Phys. Slov. **49**, 491 (1999); A. Kuzmich, I. A. Walmsley, and L. Mandel, Phys. Rev. Lett. **85**, 1349 (2000); Z.-B. Chen, J.-W. Pan, G. Hou, and Y.-D. Zhang, Phys. Rev. Lett. **88**, 040406 (2002).
 - [5] M. D. Reid and P. D. Drummond, Phys. Rev. Lett. **60**, 2731 (1988); M. D. Reid, Phys. Rev. A **40**, 913 (1989).